



Strain-specific serological response after simultaneous vaccination with PRRS MLV against

Kristensen, Charlotte Sonne; Kvisgaard, Lise Kirstine; Pawlowski, M.; Carlsen, S. H.; Hjulsager, Charlotte Kristiane; Larsen, Lars Erik

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Kristensen, C. S., Kvisgaard, L. K., Pawlowski, M., Carlsen, S. H., Hjulsager, C. K., & Larsen, L. E. (2015). *Strain-specific serological response after simultaneous vaccination with PRRS MLV against*. Abstract from International Porcine Reproductive And Respiratory Syndrome Congress, Ghent, Belgium.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Strain-specific serological response after simultaneous vaccination with PRRS MLV against PRRSV types 1 and 2: Impact on interpretation of surveillance data

C. S. Kristensen^{1*}, L.K. Kvisgaard², M. Pawlowski², S. H. Carlsen², C.K. Hjulsager², L.E. Larsen²

¹Danish Pig Research Centre, SEGES, Kjellerup, Denmark.

²Technical University of Denmark; National Veterinary Institute, Frederiksberg, Denmark

Both type 1 (subtype 1) and type 2 PRRSV are currently circulating in Denmark. In some double infected herds, the pigs are simultaneously vaccinated with PRRSV modified live vaccines (MLV) against both genotypes. There is a lack of data on the impact on serological response following simultaneous administration of PRRSV MLVs against both type 1 and type 2 PRRSV. The objective of this experimental study was to compare the level of serological response of single-vaccinated pigs with responses of double-vaccinated pigs and to compare the serological response of the two vaccination strategies following challenge with homologous and heterologous virus strains.

Sixty-six four-week-old PRRSV-negative pigs were included in the study. The pigs were purchased from a specific pathogen-free herd and tested free of a range of pathogens including PRRSV by serology at the beginning and end of the study. The pigs were housed at the experimental animal facilities at the National Veterinary Institute under appropriate biosecurity conditions. On arrival (week 0), the pigs were randomly distributed into four groups (1-4). Each group was housed in a separate room. One week after arrival (week 1), the pigs in groups 1-3 were vaccinated with either Porcilis® PRRS VET or Ingelvac® PRRS VET, or both. The last group was kept as a non-vaccinated control. Nine weeks after vaccination, all pigs were divided into three new groups and were then challenged with PRRSV type 1 (DK strain 18794), PRRSV type 2 (DK strain 19407b) or PRRSV atypical strain (strain BOR59) by the intranasal route. Blood samples were collected daily from all pigs during the first week after vaccination and challenge. In the remaining periods, samples were taken once a week from all pigs. The level of antibodies against PRRS virus was measured by type-specific ELISA and IPMA.

All pigs developed antibodies against PRRSV two to three weeks after vaccination as measured by ELISA and IPMA. The onset and level of antibodies developing in response to single and double vaccination were equal. Similarly, the development of type-specific antibodies in pigs vaccinated with the type 2 vaccine was comparable to that in pigs vaccinated with the type 1 vaccine. The profile of antibody responses following challenge with the atypical PRRSV type 1 strain BOR59 was identical to the responses of the animals challenged with the subtype 1 strain. These results indicated that simultaneously vaccination with two different PRRSV vaccines is feasible, although they also revealed that standard serological assays cannot distinguish between infections with the different type 1 subtypes. This emphasises the fact that viral detection and genomic characterisation are required for adequate surveillance of circulating PRRSV type 1 strains in Europe.

This work was partly funded by the Boehringer Ingelheim European PRRS Research Award 2014. The BOR59 isolate was kindly provided by Tomasz Stadejek, University of Warsaw.